

Anomalous finite size corrections in random field models

C. Lucibello,¹ F. Morone,² G. Parisi,³ F. Ricci-Tersenghi,³ and Tommaso Rizzo⁴

¹*Dipartimento di Fisica, "Sapienza" University of Rome, P.le A. Moro 2, I-00185, Rome, Italy*

²*Levich Institute and Physics Department, City College of New York, New York, New York 10031, USA*

³*Dipartimento di Fisica, IPCF-CNR, UOS Roma, and INFN, Sezione di Roma1,*

"Sapienza" University of Rome, P.le A. Moro 2, I-00185, Rome, Italy

⁴*IPCF-CNR, UOS Roma, "Sapienza" University of Rome, P.le A. Moro 2, I-00185, Rome, Italy*

The presence of a random magnetic field in ferromagnetic systems leads, in the broken phase, to an anomalous $O(\sqrt{1/N})$ convergence of some thermodynamic quantities to their asymptotic limits. Here we show a general method, based on the replica trick, to compute analytically the $O(\sqrt{1/N})$ finite size correction to the average free energy. We apply this method to two mean field Ising models, fully connected and random regular graphs, and compare the results to exact numerical algorithms. We argue that this behaviour is present in finite dimensional models as well.

I. INTRODUCTION

There is a large interest in understanding the behaviour of ferromagnetic systems in random magnetic fields. Also the most simple model one can conceive, the Random Field Ising Model (RFIM), has a plethora of applications to condensed matters problems [1–3], while a complete understanding of its static and dynamical properties is still lacking. Only in recent years some long-standing issues such as the presence of a glassy phase [4, 5] and the universality of the zero-temperature fixed point [6] have been partially settled.

Here we contribute to the literature on such random systems discussing the convergence properties of the average free energy to its thermodynamic limit. Finite size corrections to the free energy have been investigated by the authors in two papers [7, 8] focusing on the paramagnetic phase of disorder Ising models on diluted graphs, in an effort to characterize the free energy correction induced by loops to the Bethe mean field approximation.

Perturbative expansions around mean field approximations are not yet numerically and analytically manageable, in spite of the large amount of work done in these years [9–11]. It was suggested in Refs. [7, 8] that the study of finite size correction for infinite range systems has many points in common with the computation of loop corrections in finite dimensional problems. In order to make progresses the case of finite size correction to the average free energy was carefully studied in the case of Erdős-Rényi and random regular graphs [7, 8]. The correction were computed in the high temperature phase and they were divergent at the transition point. We show in this paper that the $O(1/N)$ corrections are overshadowed in the whole low temperature phase, because $O(\sqrt{1/N})$ corrections arise. The computation of these corrections is the goal of this paper. We present both numerical simulations and analytical computations to support this claim.

These $O(\sqrt{1/N})$ corrections do not arise from loops, but they come from disorder induced fluctuations in the weight of the two macroscopically different states. A more complex version of this phenomenon should take

place in the presence of many different equilibrium states. This phenomenon is also present in finite dimensional models, as it is suggested by some preliminary numerical we conducted. From the technical point of view the analytical computations in the replica framework are connected to the existence of multiples solutions of the mean field equations [12][13].

In the following Section we present the general analytical framework that can be employed to compute the $O(1/\sqrt{N})$ corrections to the average free energy density. Notice that the same corrections are of order $O(\sqrt{N})$ when referred to extensive quantities, therefore the indistinct use in the text of $O(1/\sqrt{N})$ and $O(\sqrt{N})$ should not confuse the reader. In Sections III and IV we apply this formalism to the fully connected and to a diluted Ising model respectively, checking numerically the consistency of the results.

II. GENERAL FORMALISM

We consider for concreteness a system of N Ising spins, $\sigma_i = \pm 1$, and external random fields. The following arguments though apply to a general class of models possessing $O(m)$ symmetry (once the average over disorder is taken). On a given graph G , the Hamiltonian of the RFIM is

$$\mathcal{H} = -J \sum_{(i,j)} \sigma_i \sigma_j - \sum_{i=1}^N h_i \sigma_i, \quad (1)$$

where the first sum is over adjacent spins, $J \geq 0$ is a ferromagnetic coupling and the the fields h_i are quenched i.i.d. random variables with zero mean and unit variance, i.e. $\mathbb{E}[h_i] = 0$ and $\mathbb{E}[h_i h_j] = \delta_{ij}$.

It takes a simple argument to show that, at least at zero temperature and for J large enough, the subleading term in N to the average energy $E(N)$ is of order $O(\sqrt{N})$. In fact in this case, for a given realization of the external fields $\{h_i\}$, the Gibbs measure is concentrated on the configuration with minimum energy, the candidates being the one with all the spins up and the one with all the

spins down. For a given graph with $M = O(N)$ edges, the energies of the two states, let us call them E_+ and E_- respectively, are given by

$$E_{\pm} = -MJ \mp \sum_{i=1}^N h_i \quad \text{for } J \gg 1. \quad (2)$$

The sum in the r.h.s. is a random variable of variance N , converging to a Gaussian variable in the thermodynamic limit. Therefore it is easy to see that for the average energy $E(N) = \mathbb{E}[\min(E_+, E_-)]$ we have

$$E(N) = -MJ - \sqrt{\frac{2}{\pi}} \sqrt{N} + o(\sqrt{N}) \quad \text{for } J \gg 1. \quad (3)$$

It turns out that the \sqrt{N} subleading behaviour we found in this limit case is present in the whole ferromagnetic region in the $J - T$ (coupling-temperature) plane.

In fact in the ferromagnetic phase the statistical weight is concentrated on two disconnected regions of the configuration space, having positive and negative magnetization respectively, separated by a free energy barrier exponentially increasing in N . While the two regions are completely equivalent in the pure ferromagnetic system, once the disordered external field is turned on their free energies start to differ of a random quantity of order $O(\sqrt{N})$. The total free energy of the system is then simply given by the minimum among the two, except for some exponentially decaying terms. Let us first define, for a given realization of the disorder, the free-energies F_+ and F_- as the ones corresponding to configurations having positive and negative magnetization respectively (for simplicity we assume N to be odd). We assume, and verify a posteriori for the RFIM, that in the ferromagnetic phase the difference between the free energies of the two states, $F_+ - F_-$, is of order $O(\sqrt{N})$. Then the average free energy of the system is given by

$$\begin{aligned} F(N) &= -\frac{1}{\beta} \mathbb{E}[\log(e^{-\beta F_+} + e^{-\beta F_-})] \\ &= \mathbb{E}[\min(F_+, F_-)] + \text{exp. vanish. terms.} \end{aligned} \quad (4)$$

From these premises it follows simply that the average free energy has an expansion in N of the form

$$F(N) = f_0 N + f_1 \sqrt{N} + o(\sqrt{N}), \quad (5)$$

where the coefficients f_0 and f_1 of the expansion are defined by

$$f_0 = \lim_{N \rightarrow +\infty} \frac{1}{N} \mathbb{E}[F_+] = \lim_{N \rightarrow +\infty} \frac{1}{N} \mathbb{E}[F_-], \quad (6)$$

$$f_1 = \lim_{N \rightarrow +\infty} \frac{-1}{\sqrt{N}} \frac{\mathbb{E}[|F_+ - F_-|]}{2}. \quad (7)$$

To compute f_1 we extend a method developed in Refs. [14] and [15–17]. At fixed N and for a given realization

of the disorder, let us call Z_+ and Z_- , the partition function constrained to the configurations with positive and negative magnetization. Considering n and m replicated systems respectively, one can easily see that for small n and m we have

$$\begin{aligned} -\frac{1}{\beta} \log \mathbb{E}[Z_+^n Z_-^m] &\sim n \mathbb{E}[F_+] + m \mathbb{E}[F_-] - \frac{n^2}{2} \beta \text{Var}(F_+) \\ &\quad - \frac{m^2}{2} \beta \text{Var}(F_-) - nm \beta \text{Covar}(F_+, F_-). \end{aligned} \quad (8)$$

Obviously, for symmetry reasons, $\mathbb{E}[F_+] = \mathbb{E}[F_-]$ and $\text{Var}[F_+] = \text{Var}[F_-]$. The free energies F_+ and F_- are jointly distributed random variables. If the limit

$$\begin{aligned} \Delta^2 &\equiv \lim_{N \rightarrow \infty} \frac{1}{2N} [\text{Var}(F_+) - \text{Covar}(F_+, F_-)] \\ &= \lim_{N \rightarrow \infty} \frac{1}{4N} \mathbb{E}[(F_+ - F_-)^2] \end{aligned} \quad (9)$$

is non-zero, the rescaled variables f_+ and f_- , defined by

$$f_{\pm} = \frac{F_{\pm} - f_0 N}{\sqrt{N}}, \quad (10)$$

become non-trivially jointly distributed Gaussian random variables for large N . In fact, while higher orders cumulants in the n and m expansion of Eq. (8) are of order $O(N)$, they give vanishing contributions to f_+ and f_- .

Since asymptotically $\frac{f_+ - f_-}{2}$ is itself a Gaussian random variable of zero mean and variance Δ^2 , it follows that the coefficient f_1 of Eqs. (5) and (7) is given by

$$f_1 = - \int \frac{dz}{\sqrt{2\pi\Delta^2}} |z| e^{-\frac{z^2}{2\Delta^2}} = -\Delta \sqrt{\frac{2}{\pi}}. \quad (11)$$

The computation of the n and m expansions in Eq. (8) can be performed using the standard replica techniques. Since it is impractical to work with the partial partition functions Z_+ and Z_- , we define the partition function Z_H of the system having an additional deterministic external field H acting on all the spins. Then we define the replicated free energy $\phi(n, m)$ as

$$\phi(n, m) = \lim_{H \rightarrow 0^+} \lim_{N \rightarrow \infty} -\frac{1}{\beta N} \log \mathbb{E}[Z_H^n Z_{-H}^m]. \quad (12)$$

This definition is completely consistent with the *l.h.s.* of Eq. (8) in the ferromagnetic region, and is even more physically sound in the paramagnetic region, allowing to treat with a unified formalism the whole phase space.

We compute $\phi(n, m)$ for integer values of n and m using the saddle point technique for $N \rightarrow \infty$. We obtain as usual a variational expression depending on a replicated order parameter and the particular structure of the field in Eq. (12) implies that we must look for a solution where the first n replicas are positively magnetized while the remaining m replicas are negatively magnetized. Then we

make an analytical continuation of the solution to real value of n and m . By definition (12) the 0 - th order term of the expansion is zero. Since we are interested only in the quadratic terms of the $\phi(n, m)$ expansion, and since the $\phi(n, m)$ is variational in some order parameter, our calculations will involve only the order parameter computed at $n = m = 0$ and we do not have to consider its n and m dependence. As always happens with calculations involving replicas, the exchange of the n and N limits will be done without particular care.

It is interesting to note that the $O(\sqrt{N})$ corrections will be obtained evaluating expression (8) at the leading $O(N)$ order, that is by taking the saddle point, while the more common $O(1)$ corrections are typically controlled by Gaussian fluctuations around the saddle point.

In the following Sections we will explicitly compute $\phi(n, m)$ given by Eq. (12) in two mean field models.

III. FULLY CONNECTED

The first case we consider is that of the RFIM on the fully connected graph. Here the first sum in the Hamiltonian (1) runs over $N(N-1)/2$ edges, and the coupling J has to be rescaled by a factor N^{-1} in order to obtain an extensive thermodynamic behaviour. It is then easy to compute the replicated free energy of the system, defined in Eq. (12), through standard Hubbard-Stratonovich transformation is given by

$$\phi_{FC}(n, m) = \min_{x \geq 0} \left\{ \frac{1}{2} J(n+m)x^2 - \frac{1}{\beta} \log \mathbb{E}_h \left[(2 \cosh \beta(Jx+h))^n (2 \cosh \beta(-Jx+h))^m \right] \right\}. \quad (13)$$

Notice that the order parameter x , the magnetization of the up-state, enters in the last term of Eq. (13) both with a plus and a minus sign. As a technical note, in Eq. (13) the minimization condition does not ever turn to a maximization when the number of replicas is small, as it happens in glassy models [18], since the dimension of the order parameter in the full replica space is $n+m$ and is always greater than zero in our calculations.

To obtain the $O(\sqrt{N})$ contribution to the free energy we have to compute the small n and m expansion of $\phi_{FC}(n, m)$. As already discussed at the end of the previous section, only the $n = m = 0$ saddle point solution, given by the non-negative solution of

$$x = \mathbb{E}_h \tanh \beta(Jx+h), \quad (14)$$

appears in the second order expansion of the replicated action. Therefore the coefficient Δ , defined in Eq. (9) and characterizing the $O(\sqrt{N})$ correction though (11), is

given by

$$\Delta_{FC}^2 = \frac{1}{2\beta^2} \left\{ \mathbb{E}_h [(\log \cosh \beta(Jx+h))^2] - \mathbb{E}_h [\log \cosh \beta(Jx+h) \log \cosh \beta(Jx-h)] \right\}. \quad (15)$$

Last expression can be easily computed for any value of J and β . In order to compare the analytical prediction with exact results, it is easier to consider the zero temperature limit of (15). This is given by

$$\lim_{T \rightarrow 0} \Delta_{FC}^2 = \frac{1}{4} \mathbb{E}_h (|Jx+h| - |Jx-h|)^2 \quad (16)$$

with x non-negative solution of

$$x = \mathbb{E}_h \text{sign}(Jx+h). \quad (17)$$

We focus on the zero temperature limit for two reasons: the zero temperature fixed point is the one controlling the flow of the renormalization group also starting from finite temperature [19]; the ground state of the RFIM can be computed efficiently using exact numerical algorithms.

We implemented an exact and very efficient algorithm to compute the ground state of a fully connected RFIM, that takes advantage of the topological equivalence of all the spins. In fact it is easy to realize that among all the configurations having a certain total magnetization $M = \sum_i \sigma_i$, the one with lowest energy is the one where only the first $\frac{N-M}{2}$ spins with the lowest external field are down.

Therefore, once the spins are sorted according to their external fields (an operation of time complexity $\Theta(N \log N)$), we have to look only to these $N+1$ configurations characterized by $M = -N, -N+2, \dots, N$ to find the ground state (an $\Theta(N)$ operation). We have thus produced an algorithm of time complexity $\Theta(N \log N)$ and with $\Theta(N)$ memory requirements. This is a great improvement over the min-cut algorithm that we use on diluted graphs (as we shall explain in next Section), that has time and memory complexity $\Theta(N^3)$ for fully connected graphs. With this algorithm we were able to perform highly precise averages of systems up to $\sim 10^6$ spins. We then subtract the leading order (in N) term $f_0 N$, that can be computed exactly, to obtain the numerical estimate of the coefficient f_1 up to subleading finite size effects. In Fig. 1 we show the perfect agreement between the results of our exact algorithm and the analytical prediction $f_1 = -\Delta \sqrt{\frac{2}{\pi}}$, with Δ given by Eq. (16), for the RFIM at $T = 0$.

IV. RANDOM REGULAR GRAPHS

The analytical computation of the $O(\sqrt{N})$ correction to the average free energy in diluted models is slightly more involved. Here we focus on the random regular graph (RRG) ensemble. Each element of the ensemble is




FIG. 1. The numerical estimates, obtained through the algorithm described in the text, of the coefficient of the $O(\sqrt{N})$ correction to the average free energy in the RFIM at zero temperature on the Fully Connected Graph, for many sizes of the system (*orange lines*). The numerical curves extrapolate to the analytical prediction of Eqs. (11) and (16) for large N (*black line*). The dashed black line is the asymptotic value $\sqrt{\frac{2}{\pi}}$.

chosen uniformly at random among all the graphs where each node has fixed connectivity z . The RRG and other diluted graph ensembles have the property of being locally tree-like, in the sense that each finite neighborhood of a randomly chosen node is with high probability a tree in the large N limit, and the density of finite loops goes to zero [20]. This property allows for the analytical solution of such models, at least at the leading order in

Under replica symmetry assumptions the order parameter $\rho(\sigma)$ can be parametrized in the form

$$\rho(\sigma) = \int \frac{dP(u^+, u^-)}{(2 \cosh \beta u^+)^n (2 \cosh \beta u^-)^m} e^{\beta(u^+ \sum_{a=1}^n \sigma^a + u^- \sum_{a=n+1}^m \sigma^a)}. \quad (21)$$

In the small n and m limit the minimum condition gives

$$P(u^+, u^-) = \mathbb{E}_h \int \prod_{k=1}^{z-1} dP(u_k^+, u_k^-) \delta\left(u^+ - g\left(h + \sum_k u_k^+\right)\right) \delta\left(u^- - g\left(h + \sum_k u_k^-\right)\right), \quad (22)$$

where the function $g(x)$ is the usual cavity iteration rule $g(x) = \frac{1}{\beta} \operatorname{atanh}(\tanh(\beta J) \tanh(\beta x))$. The vanishing auxiliary external field H of Eq. (12) selects the solution of Eq. (22), supposed to be unique, such that $\overline{u^+} \geq 0 \geq \overline{u^-}$ (denoting with $\overline{\cdot}$ the expectation over $P(u^+, u^-)$). The algorithmic equivalent of the distributional equation (22) is nothing else than the standard Belief Propagation algorithm applied two times with opposite initializations on the same sample.

Expanding the replicated free energy $\phi_{RRG}(n, m)$ to

N .

Following the lead of Ref. [17], we apply the replica technique to Eq. (12), to obtain the replicated free energy

$$\phi_{RRG}(n, m) = \min_{\rho} \left\{ \frac{z}{2} \phi_{edge}[\rho] - (z-1) \phi_{site}[\rho] \right\}, \quad (18)$$

where

$$\begin{aligned} \phi_{edge}[\rho] &= \log \left[\mathbb{E}_h \sum_{\sigma_1, \sigma_2} \rho^{z-1}(\sigma_1) \right. \\ &\quad \left. \times e^{\beta(h_1 \sum_a \sigma_1^a + J \sum_a \sigma_1^a \sigma_2^a + h_2 \sum_a \sigma_2^a)} \rho^{z-1}(\sigma_2) \right]. \end{aligned} \quad (19)$$

and

$$\phi_{site}[\rho] = \log \left[\mathbb{E}_h \sum_{\sigma} e^{\beta h \sum_{a=1}^{n+m} \sigma^a} \rho^z(\sigma) \right] \quad (20)$$

A similar expression, for the replicated free energy of spin glasses on RRG, was derived in Ref. [21] (see also Eq. (7) of Ref. [22] for a more general formulation) and relies on the hypothesis that the graph contains few short loops.

Eq. (18) is a straightforward generalization of these results, the difference being that the replicas are divided into two blocks of size n and m respectively. The minimization condition in Eq. (18) is imposed over all the functions $\rho(\sigma) = \rho(\sigma^1, \dots, \sigma^{n+m})$ of a $n+m$ replicated Ising spin, taking particular attention to constrain the first n and last m replicas to be in the up and down state respectively. Here, as in the fully connected case, the dimension of the order parameter is always positive, since it is 2^{n+m} . Therefore we do not incur in a common peculiarity of replica calculations, the exchange between minimization and maximization conditions [18].

the second order in n and m we can then derive the $O(\sqrt{N})$ coefficient of the free energy according to Eqs. (8) and (11). We note that only the fixed point distribution $P(u^+, u^-)$ computed at $n = m = 0$, that is the solution of (22), contributes to the quadratic order of $\phi(n, m)$. The coefficient Δ appearing in $f_1 = -\Delta \sqrt{2/\pi}$

than reads

$$\Delta_{RRG}^2 = \frac{1}{2\beta^2} \left\{ \frac{z}{2} (\mathbb{E}[A_+^2] - \mathbb{E}[A_+A_-]) - (z-1) (\mathbb{E}[B_+^2] - \mathbb{E}[B_+B_-]) \right\}. \quad (23)$$

The terms A_\pm and B_\pm stem from the edge and site replicated free energies (ϕ_{site} and ϕ_{edge}) respectively. The expectations $\mathbb{E}[\cdot]$ are over both the distribution of the external random field and of the cavity fields. The site terms B_+ and B_- are defined by

$$B_\pm = \log \frac{2 \cosh \beta(h + \sum_{k=1}^z u_k^\pm)}{\prod_{k=1}^z 2 \cosh \beta u_k^\pm}. \quad (24)$$

Here the fields u_k^+ and u_k^- are distributed according to $P(u_k^+, u_k^-)$, solution of Eq. (22), and h is distributed as the external random fields. The edge terms A_+ and A_- , appearing in Eq. (23), are defined by

$$A_\pm = \log \sum_{\sigma_1, \sigma_2} \frac{\exp \beta(h_1^\pm \sigma_1 + J \sigma_1 \sigma_2 + h_2^\pm \sigma_2)}{\prod_{k=1}^{z-1} 4 \cosh \beta u_{1k}^\pm \cosh \beta u_{2k}^\pm} \quad (25)$$

The random field h_1^+ is distributed as $h + \sum_{k=1}^{z-1} u_{1k}^+$, where h is an external random fields, and analogous definitions follows for the other cavity fields.

We notice that only in the terms $\mathbb{E}[A_+A_-]$ and $\mathbb{E}[B_+B_-]$ of Eq. (23) the full joint distribution $P(u^+, u^-)$ is needed, not only its marginals.

The computation of Δ^2 , and therefore of the analytical finite size correction f_1 , to a high level of precision through Eq. (23), is a computationally easy task. We solved numerically the fixed point condition Eq. (22) through a population dynamic algorithm. In this case each element of the population is a couple of cavity messages, u^+ and u^- , each of them encountering the same external random fields h during the iterations of the algorithm. As initial condition, in each couple the message u^+ is set to a high positive value, while the message u^- is set to a low negative value.

In the paramagnetic phase the stable solution of Eq. (22) takes the trivial form $P(u^+, u^-) = P(u^+) \delta(u^+ - u^-)$. The $O(\sqrt{N})$ finite size correction is thus zero. In the ferromagnetic phase instead, the messages u^+ and u^- become non-trivially correlated.

We computed with the population dynamics algorithm the solution of the fixed point Eq. (22) at temperature $T = 0$, for many values of the coupling J and for connectivity $z = 4$. The expectations we find in the expression of Δ^2 given Eq. (23) are then computed sampling from the population.

As in the case of the fully connected model, it is easier to verify the analytical predictions working at zero temperature, since the free energy and the energy coincides and the ground state of the system can be obtained through exact polynomial algorithms.




FIG. 2. The numerical estimates, obtained through the min-cut algorithm, of the coefficient of the $O(\sqrt{N})$ correction to the average free energy in the RFIM on large temperature on Random Regular Graphs of connectivity $z = 4$, for many sizes of the systems (orange lines). The numerical curves (see Eq. 26 for their definition) extrapolate to the analytical prediction of Eqs. (11) and (23) for large N (black line). The dashed black line is the asymptotic value $\sqrt{2/\pi}$.

Therefore the analytical result is compared with an exact numerical algorithm that exploit the equivalence between the problem of finding the ground state of the RFIM on an arbitrary graph and the minimum cut optimization problem [23]. We used the implementation of the Goldberg-Tarjan's preflow push-relabel algorithm provided by the open source Lemon Graph Library [24], whose worst case complexity is $\Theta(N^{\frac{5}{2}})$ for instances of the RRG ensemble. The numerical estimate of the $O(\sqrt{N})$ coefficient is obtained from a linear combinations of the free energy of systems of different sizes, and it is given by

$$\tilde{f}_1(N) = \frac{\mathbb{E}[F(2N) - 2F(N)]}{c\sqrt{N}} \quad (26)$$

with $c = \frac{1}{\sqrt{2}} - 1$. The value of $\tilde{f}_1(N)$, obtained averaging the minimum cut results over many samples of the system, should converge for large N to the analytical value computed through the population dynamic algorithm applied to Eq. (23). The data plotted in Fig. 2 show a very good agreement between experiments and predictions, although the convergence is slow due to the presence of subleading $O(\frac{1}{\sqrt{N}})$ finite size effects.

Moreover the analytic value of f_1 , when plotted as a function of the absolute magnetization of the system, $m \equiv \lim_{N \rightarrow \infty} N^{-1} \mathbb{E}[|\sum_i < \sigma_i >|]$, see Figure 3, shows a weak dependence from the connectivity of the system and an almost linear behaviour. A simple estimate of f_1 on both the fully connected graph and RRGs is then given by $f_1 \approx -\sqrt{2/\pi} m$.




FIG. 3. The coefficient of the $O(\sqrt{N})$ correction to the average free energy versus the average absolute magnetization m in the RFIM at zero temperature on Random Regular Graphs of different connectivities, as computed solving Eqs. (22) and (23) with a population dynamic algorithm. The black line shows the same quantity for the Fully Connected model.

V. CONCLUSIONS

We are arguing for the existence of an anomalous $O(1/\sqrt{N})$ subleading correction to the thermodynamic average free energy f_0 of systems with a zero-mean external random field. This correction is limited to the ferromagnetic phase, since it is caused by the difference of free energy among the two pure states. If one is interested only in asymptotic quantities such as f_0 , the slowing down of the convergence can be avoided in numerical simulations such as Markov Chain Monte Carlo methods choosing an initial condition uncorrelated to the realization of the disorder (as it is indeed usually done), such that the dynamics gets trapped with equal probability in the state with lowest free energy or in the other one. On the other hand estimations of some observables on systems of finite size through exact algorithms, such as the minimum cut algorithm, are bound to follow a behaviour of the type $a(N) \sim a_0 + \frac{a_1}{\sqrt{N}}$, therefore the convergence is much slower than the usual $O(\frac{1}{N})$ behaviour one finds in the paramagnetic phase or in pure systems. Constraining

the sampling an observable to a state symmetrically fluctuating around its mean asymptotic value is in this sense more convenient than sampling from the pure state with greater statistical weight.

Using a formalism inspired by some recent works [14–17] we presented a general framework to compute the coefficient f_1 of the $O(\sqrt{N})$ term in the average free energy. The computation has been carried out, using a variant of the replica trick, in two solvable mean field systems, the RFIM on the fully connected graph and on the RRG ensemble. The analytical results obtained, Eqs. (11)(15)(23), are found to be in strong agreement with the numerical simulations.

A different but equivalent approach to the problem, based on a peculiar replica symmetry breaking scheme, can be taken using the techniques of Refs. [12, 13]. Instead of computing the large deviation function $\phi(n, m)$, where n and m replicas are constrained to be in the up and down state respectively, and expanding to small values of n and m , the same conclusions can be obtained summing over all the saddle point contributions obtained from the partitioning of a the replicas in the two sets. We preferred the approach of Refs. [14–17] because it is conceptually more clear and analytically less involving (although they share many similarities).

While the formalism we have developed in Section II is completely general, the exact computation of the coefficient Δ^2 (thus of f_1) can be achieved only in mean field models. In finite dimension one has to resort to a perturbative diagrammatic expansion of a replicated field theory. We did not take this path, but our numerical simulations with a min-cut algorithm, using the same procedure described in Section IV for the RRG ensemble, show, qualitatively and also quantitatively, the scenario depicted in Fig. 2. Summing it up, as a general feature of models with zero-mean random external field, the average free energy density as a first finite size correction of order $O(\frac{1}{N})$ in the paramagnetic phase, and of order $O(\frac{1}{\sqrt{N}})$ in the ferromagnetic phase.

The research leading to these results has received funding from the European Research Council under the European Union’s Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement No. 247328 and from the Italian Research Minister through the FIRB project No. RBFR086NN1.

-
- [1] S. Fishman and A. Aharony, Journal of Physics C: Solid State Physics **12**, L729 (1979).
 [2] E. Dagotto, Science **309**, 257 (2005).
 [3] R. Vink, K. Binder, and H. Löwen, Physical Review Letters **97**, 230603 (2006).
 [4] F. Krzakala, F. Ricci-Tersenghi, D. Sherrington, and L. Zdeborová, Journal of Physics A: Mathematical and Theoretical **44**, 042008 (2011), arXiv:1008.4497.
 [5] F. Krzakala, F. Ricci-Tersenghi, and L. Zdeborová, Physical Review Letters **104**, 207208 (2010), arXiv:0911.1551.
 [6] N. G. Fytas and V. Martin-Mayor, Physical Review Letters **110**, 227201 (2013).
 [7] U. Ferrari, C. Lucibello, F. Morone, G. Parisi, F. Ricci-Tersenghi, and T. Rizzo, Physical Review B **88** (2013), 10.1103/PhysRevB.88.184201.
 [8] C. Lucibello, F. Morone, G. Parisi, F. Ricci-Tersenghi, and T. Rizzo, ArXiv e-prints, 1 (2014), arXiv:1403.6049.

- [9] A. Montanari and T. Rizzo, *Physical Review B* **79** (2009), 10.1103/PhysRevB.79.134205, *Journal of Statistical Mechanics: Theory and Experiment* **2005**, P050108 (2005), arXiv:0506769v1 [cond-mat].
- [10] M. Chertkov and V. Y. Chernyak, *Physical Review E* **73** (2006), 10.1103/PhysRevE.73.065102, arXiv:0601487v2 [cond-mat].
- [11] G. Parisi and F. Slanina, *Journal of Statistical Mechanics: Theory and Experiment* **2006**, P02003 (2006), arXiv:0512529v1 [cond-mat].
- [12] V. Dotsenko and M. Mézard, *Journal of Physics A: Mathematical and General* **30**, 3363 (1997), arXiv:9611017 [cond-mat].
- [13] V. Dotsenko, *Journal of Statistical Mechanics: Theory and Experiment* **2006**, P060371 (2006), arXiv:0603713 [cond-mat].
- [14] T. Aspelmeier, M. A. Moore, and A. P. Young, *Physical Review Letters* **90**, 127202 (2003).
- [15] G. Parisi and T. Rizzo, *Physical Review Letters* **101** (2008), 10.1103/PhysRevLett.101.17205, arXiv:0706.1180.
- [16] G. Parisi and T. Rizzo, *Physical Review B* **79** (2009), 10.1103/PhysRevB.79.134205, arXiv:0810.24v2.
- [17] G. Parisi and T. Rizzo, *Journal of Physics A: Mathematical and Theoretical* **43** (2010), 10.1088/0305-4470/43/10/105501, arXiv:0910.4553.
- [18] G. Parisi, M. Mézard, and M. A. Virasoro, *Spin glass theory and beyond* (World Scientific Singapore, 1987).
- [19] A. Bray and M. A. Moore, *Journal of Physics C: Solid State Physics* **18**, L927 (1985).
- [20] A. Montanari and M. Mézard, *Information, Physics and Computation* (Oxford Univ. Press, 2009).
- [21] Y. Y. Goldschmidt and C. De Dominicis, *Physica* **2006**, 206041 (1990).
- [22] T. Rizzo, A. Lage-Castellanos, R. Mulet, and F. Ricci-Tersenghi, *Journal of Statistical Physics* **139**, 375 (2010), arXiv:0906.2695.
- [23] J. Picard and H. D. Ratliff, *Networks* **5**, 357 (1975).
- [24] B. Dezs, A. Jüttner, and P. Kovács, *Electronic Notes in Theoretical Computer Science* **264**, 23 (2011).