



LETTER

One-loop topological expansion for spin glasses in the large connectivity limit

To cite this article: Maria Chiara Angelini *et al* 2018 *EPL* **121** 27001

View the [article online](#) for updates and enhancements.

One-loop topological expansion for spin glasses in the large connectivity limit

MARIA CHIARA ANGELINI¹, GIORGIO PARISI^{1,2,3} and FEDERICO RICCI-TERSENGHI^{1,2,3}

¹ *Dipartimento di Fisica, Sapienza Università di Roma - Piazzale A. Moro 2, I-00185, Rome, Italy*

² *Nanotec-CNR, UOS Rome, Sapienza Università di Roma - Piazzale A. Moro 2, I-00185, Rome, Italy*

³ *INFN-Sezione di Roma 1 - Piazzale A. Moro 2, 00185, Rome*

received 23 October 2017; accepted in final form 23 February 2018
published online 16 March 2018

PACS 75.10.Nr – Spin-glass and other random models
PACS 05.10.Cc – Renormalization group methods

Abstract – We apply for the first time a new one-loop topological expansion around the Bethe solution to the spin-glass model with a field in the high connectivity limit, following the methodological scheme proposed in a recent work. The results are completely equivalent to the well-known ones, found by standard field-theoretical expansion around the fully connected model (Bray and Roberts 1980, and following works). However this method has the advantage that the starting point is the original Hamiltonian of the model, with no need to define an associated field theory, nor to know the initial values of the couplings, and the computations have a clear and simple physical meaning. Moreover this new method can also be applied in the case of zero temperature, when the Bethe model has a transition in field, contrary to the fully connected model that is always in the spin-glass phase. Sharing with finite-dimensional model the finite connectivity properties, the Bethe lattice is clearly a better starting point for an expansion with respect to the fully connected model. The present work is a first step towards the generalization of this new expansion to more difficult and interesting cases as the zero-temperature limit, where the expansion could lead to different results with respect to the standard one.

Copyright © EPLA, 2018

Introduction. – Spin glasses (SGs) are models whose mean-field (MF) version [1] undergoes a phase transition, crossing a critical line in the temperature-field (T, h) -plane. The solution of the MF problem sees the introduction of replicas of the original system as a mathematical trick to perform computations. The resulting Hamiltonian is symmetric under replica exchanges. However, quite surprising, one finds that in the low-temperature spin-glass phase the replica symmetry is broken. While the MF behavior of the model is completely under control [2], also from a rigorous viewpoint [3], we still do not have a confirmed theory for the finite-dimensional version. In particular, there is not agreement both on the upper and lower critical dimension, looking at theoretical, numerical and experimental data [4–12].

The project to perform a renormalization group (RG) analysis is an old one. The spin-glass transition in zero-field was already studied within the RG by Harris *et al.* [13], and in field by Bray and Roberts [14], limiting at the sector associated with the critical eigenvalue, the so called *replicon*. Their one-loop analysis was then repeated adding the other sectors, longitudinal and anomalous ones,

in refs. [5,15–17]. In a recent work also the two-loop computation in a field has been performed [7], suggesting the possibility of a non-perturbative fixed point.

These works are expansions around the fully connected (FC) mean-field model. They start studying the symmetric phase, approaching the transition from the high-temperature side, where replica symmetry holds. Thus, the replica symmetric Lagrangian is written, that in its most complete version has three bare masses and eight cubic couplings involving the replica fields, which correspond to all the possible invariants under the replica symmetry. At this point one can perform a renormalization à la Wilson, integrating the degrees of freedom over an infinitesimal momentum shell, extracting the leading, one-loop, order approximation in $\epsilon = 6 - d$. Although the scheme is clear, the computation is highly technical also from the algebraic viewpoint.

Recently a new loop expansion around the mean-field Bethe solution was proposed in ref. [18]. The new method can be applied to each model that is well defined on a Bethe lattice. In this paper, we apply for the first time this new expansion to the SG in a field. We restrict ourselves

to the limit of high connectivity $z \rightarrow \infty$ to perform computations analytically. We compute the 1st-order correction, and we show that in the $T > 0$ region this new expansion is completely equivalent to the field-theoretical one, recovering the results of Bray and Roberts [14]. However it has the advantage that the starting point is the original Hamiltonian of the model, with no need to define an associated field theory, nor to know the initial values of the couplings, and the computations have a clear and simple physical meaning: while in standard field theory Feynmann diagrams have no special meaning, here the important diagrams have a geometrical interpretation. Moreover, the expansion is around Bethe lattice that has finite connectivity, an important characteristic shared with finite-dimensional systems. Even if in this work we obtain the same results as standard RG around the fully connected model, we will discuss the differences that could arise in the two methods in particular situations. This work is first of all a verification of the correctness of the method proposed in ref. [18] and it is a first step towards the generalization of this new expansion to more complicated cases: finite small connectivity and zero temperature.

Expanding around the Bethe lattice solution. –

In this section we just recap the results of ref. [18], while in following sections we will apply for the first time these results to the SG model in a field. Starting from a D -dimensional system, the M -layer construction of ref. [18] consists of taking M copies of the original model and rewire them. In the $M \rightarrow \infty$ limit, the rewiring procedure leads to the Bethe lattice. One can then expand the observables in powers of $\frac{1}{M}$. In particular, we will focus our attention on the correlation functions, connected over the disorder, let us name them G . The $\frac{1}{M}$ expansion results in a topological expansion in the number of loops. At order $\frac{1}{M}$, the correlation function between the origin and a point d in the D -dimensional model results to be: $G(d) = \sum_{L=1}^{\infty} \mathcal{B}(d, L) g^B(L)$, where $\mathcal{B}(d, L)$ is the number of non-backtracking walks that go from the origin to the point d in the original D -dimensional model and $g^B(L)$ is the correlation between points at distance L on a Bethe lattice.

At order $\frac{1}{M^2}$, the correlation function on the original system receive a leading contribution that is the product of the so-called *line-connected* observable $g^{lc}(\mathcal{L})$ computed on a Bethe lattice, in which it has been manually injected a loop \mathcal{L} of the type in fig. 1, multiplied by the number of such a structure \mathcal{L} present on the original model. The line-connected observable is just the observable computed on a Bethe lattice with the loop minus the observable computed on the two paths $L_0, L_1 + 1, L_3$ and $L_0, L_2 + 1, L_3$ considered as independent.

The quantities $g^B(L)$ and $g^{lc}(\mathcal{L})$ are model dependent. In the following we will compute them for the spin glass in a field. To make the computation analytically feasible, we will compute things on a Bethe lattice in the high




Fig. 1: Spatial loop \mathcal{L} that gives the first correction to the bare correlation functions in the expansion around the Bethe solution.

connectivity limit, at temperature $T > 0$. However things can be computed in finite connectivity and even at $T = 0$ using numerically the Belief Propagation equations. This will be the subject of a subsequent paper.

We just want to recall that one could perform the same M -layer construction around the fully connected model instead of the Bethe lattice. In the former approach, the leading divergences at each order are exactly given by the corresponding terms in the loop expansion of the continuum field theory. In ref. [18] it is claimed that, if the critical behaviour of the fully connected model and of the Bethe lattice model is the same, then also the two expansions will lead to the same results. We will see that this is exactly the case for the SG with field at $T > 0$.

Model and definitions. – To be concrete, we are interested in the SG model in a field, that has the following Hamiltonian:

$$H(\{\sigma\}) = - \sum_{i=1, N} \sigma_i h_i^R - \sum_{ij \in E} \sigma_i \sigma_j J_{ij},$$

where E is the set of the edges of the lattice. On this model we should compute the quantities $g^B(L)$ and $g^{lc}(\mathcal{L})$ introduced in the previous section.

We will consider the model on a Bethe lattice (for definiteness on a random z -regular lattice) in the large connectivity limit, *i.e.*, z large: we will keep the leading terms and we will neglect the $1/z$ corrections. Only at the end we will perform the limit $z \rightarrow \infty$. In this limit, computations are easier and the model has the same properties of the Sherrington-Kirkpatrick one [1]. The procedure of first computing the results for finite z in the thermodynamic limit ($N \rightarrow \infty$) and later send z to infinity makes the physical approach much clearer.

The couplings are i.i.d. random variables extracted from a distribution with the following properties: $\overline{J_{ij}} = 0$, $\overline{J_{ij}^2} = 1/z$. Higher-order moments are irrelevant in the $z \rightarrow \infty$ limit. We have indicated by h_i^R the field on the site i . It can be either a local random field extracted from a given distribution or a spatially uniform field. The physics is equivalent in the two cases. For simplicity here we consider the case where the fields h_i^R are Gaussian variables with zero average and finite variance v_h (see footnote ¹).

¹The attentive reader could notice that the variance v_h never appears in the following. This is not because things do not depend on v_h , but because the dependence is hidden in the definitions of the magnetizations: $m^2 \equiv \langle \sigma \rangle^2$ and higher moments will implicitly depend on v_h .

$$\overline{Z^n} = \overline{\sum_{\{\sigma\}} e^{\beta \sum_a \sum_i \sigma_i^a h_i^R} e^{\beta \sum_a \sum_{ij} \sigma_i^a \sigma_j^a J_{ij}}} \simeq \overline{\sum_{\{\sigma\}} e^{\beta \sum_a \sum_i \sigma_i^a h_i^R} \prod_{ij} \left(1 + \beta \sum_a \sigma_i^a \sigma_j^a J_{ij} + \frac{\beta^2}{2} \sum_{a,b} \sigma_i^a \sigma_j^b \sigma_j^a \sigma_i^b J_{ij}^2 \right)}. \quad (1)$$

With standard notation, we indicate with $\langle \cdot \rangle$ the thermal average and with $\overline{\cdot}$ the average over the disorder (random couplings and fields).

In the thermodynamic limit we will compute different kinds of correlation functions first between point at distance L on a standard Bethe lattice; this will lead to the *bare* propagator and we will compute its exact expression with two different methods in the high-temperature region: the replica method and the cavity method. Then we will compute the first correction to this result due to the presence of one spatial loop. The limit $z \rightarrow \infty$ allows us to compute all the quantities analytically. For finite connectivity, one could compute everything numerically using Belief Propagation as usually done on Bethe lattices.

The replica computation of line-correlations. – In the high connectivity limit (that corresponds to small couplings limit) we can expand the replicated partition function as

see eq. (1) above

The neglected terms give sub-leading contributions for large z .

As usual in SG computations, $a, b, c, \dots \in [1, n]$ indicate the replica index, where the replicas are independent copies of the system with the same disorder realization.

A Bethe random regular graphs in the $N \rightarrow \infty$ limit becomes locally loop-less. The distance on the graph between two generic points (*i.e.*, the length of the shortest path between them) is of order $\log(N)$. We are interested in the computation of the correlation functions of spins that are on points at a distance L between them, in the limit where N goes to infinity at fixed L . In this limit with probability one there is a unique path (of finite length) connecting them so the computation can be done on a single line.

In general we will be interested in correlation functions that are connected with respect to the disorder. At this end it is convenient to compute

$$G_{a,b,c,d}(L) \equiv \overline{\langle \sigma_0^a \sigma_0^b \sigma_L^c \sigma_L^d \rangle} - \overline{\langle \sigma_0 \rangle}^2 \cdot \overline{\langle \sigma_L \rangle}^2, \quad (2)$$

from which we can extract connected and disconnected (with respect to thermal average) correlation functions. In the following, for simplicity of notation, we will indicate with $\overline{\langle \cdot \rangle}^c$ the correlations connected with respect to the disorder: $\overline{\langle \sigma_0^a \sigma_0^b \sigma_d^c \sigma_d^d \rangle}^c \equiv \overline{\langle \sigma_0^a \sigma_0^b \sigma_d^c \sigma_d^d \rangle} - \overline{\langle \sigma_0 \rangle}^2 \cdot \overline{\langle \sigma_d \rangle}^2$. We define the matrix $T \in \mathbb{R}^{n(n-1) \times n(n-1)}$ such as $T_{ab,cd}(i) \equiv \frac{\beta^2}{2} \overline{\langle \sigma_i^a \sigma_i^b \sigma_i^c \sigma_i^d \rangle}$. We define T only for $a \neq b$,

$c \neq d$, following what is usually done for the matrix $Q_{ab}(i) = \langle \sigma_i^a \sigma_i^b \rangle$ in replica calculations². Remembering that $\overline{J_{ij}} = 0$, $\overline{J_{ij}^2} = \frac{1}{z}$, from eq. (1) we find that

$$G_{a,b,c,d}(L) = \frac{2}{\beta^2 z^L} \left[\prod_{i=0}^L T(i) \right]_{ab,cd}. \quad (3)$$

Please notice that all the $i \in [0, L]$ are present in eq. (3). In fact on a Bethe lattice, two spins are linked just by a path. If the link between two spins is cut, the spins become disconnected. This means that if a coupling is 0, all the correlation functions between the two spins linked by that coupling are zero. This automatically implies that a correlation function should be proportional to the product of all the couplings on the path between the two spins.

From eq. (3), we need to compute powers of the matrix T . It is easy to show that

$$T_{ab,cd}(i) = \frac{\beta^2}{2} \overline{\langle \sigma_i^a \sigma_i^b \sigma_i^c \sigma_i^d \rangle} = \frac{\beta^2}{2} \cdot \begin{cases} 1, & \text{if } a = c, b = d \text{ or if } a = d, b = c, \\ m_2, & \text{if } a = c \text{ or } b = d \text{ or } a = d \text{ or } b = c, \\ m_4, & \text{if } a \neq b \neq c \neq d \end{cases} \quad (4)$$

with $m_2 = \overline{\langle \sigma_i \rangle}^2$ and $m_4 = \overline{\langle \sigma_i \rangle}^4$. Let us just mention that for eq. (3) to hold, T should be defined as the so-called ‘‘cavity’’ average: the average over the rest of the system with the exception of the neighbouring spins on the considered line. However, in the large z limit, cavity averages are equal to standard averages (see also the Supplementary Material [Supplementarymaterial.pdf](#) (SM)).

Equation (4) can be written in the form

$$T_{ab,cd} = \frac{\beta^2}{2} \cdot [m_4 + (m_2 - m_4)(\delta_{ad} + \delta_{bc} + \delta_{bd} + \delta_{ac}) + (1 - 2m_2 + m_4)(\delta_{ac}\delta_{bd} + \delta_{ad}\delta_{cb})]. \quad (5)$$

In order to compute correlation functions, we need to compute powers of T . For this reason, we proceed to the diagonalization of T . The whole calculation is explained in the SM. Here we just sketch the main steps and state the final result. First of all we look for eigenvalues and eigenvectors of $T_{ab,cd}$, of the form: $\sum_{cd} T_{ab,cd} \psi_{cd} = \lambda \psi_{ab}$.

²Please be careful to not confuse T with a tensor. We could use two superindices $i, j \in [1, n(n-1)]$ instead of the couples ab, cd . However we choose this notation because it will be useful to define the different types of correlation functions in the following.

Because of the symmetry of the matrix T under permutations of the replica indices, we know that there are three symmetry classes of eigenvectors (in an analogous way to what one does when looking to the stability of the Sherrington-Kirkpatrick solution for the FC model [19]) and three associated eigenvalues:

In the limit $n \rightarrow 0$ the first two eigenvalues (longitudinal and anomalous) are $\lambda_{L/A} = \beta^2(3m_4 - 4m_2 + 1)$, while the third one (replicon) is $\lambda_R = \beta^2(1 - 2m_2 + m_4)$.

We just want to point out that the eigenvalues are not the same ones as the usual “replicon, anomalous, longitudinal” eigenvalues that comes out from the diagonalization of the Hessian in ref. [19], but we called them in the same way because they identify the same sub-spaces with the same replica symmetries. In particular at the spin-glass transition the usual replicon goes to zero, while λ_R as defined in this paper goes to $\lambda_R = 1$ leading to the divergence of the spin-glass susceptibility (see the following section).

At this point, we construct the projectors on the sub-spaces of the eigenvectors and write T as a combination of the projectors. In this representation, it is easy to compute powers of T . A special care should be taken in performing the limit $n \rightarrow 0$, because of the degeneration of λ_L and λ_A . The final result is

$$T^L(n=0) = \frac{\beta^2}{2} L \lambda_{L/A}^{L-1} (3m_4 - 2m_2) R + \lambda_{L/A}^L \left(-\frac{R}{2} - Q \right) + \lambda_R^L \left(\frac{R}{2} + Q + P \right),$$

where we defined the matrices $R_{ab,cd} = 1$, $Q_{ab,cd} = \frac{1}{4}[\delta_{ac} + \delta_{ad} + \delta_{bc} + \delta_{bd}]$, $P_{ab,cd} = \frac{1}{2}[\delta_{ac} \cdot \delta_{bd} + \delta_{bc} \cdot \delta_{ad}]$.

For a spin-glass model, for each realization of the system different correlation functions can be defined. Because of the symmetry of the coupling distribution, the average over the realizations of all the “linear” correlations will be zero, and the relevant ones will be the squared correlations. We will define three main correlations averaged over the thermal noise:

- The total correlation: $\langle \sigma_0 \sigma_L \rangle$.
- The disconnected correlation: $\langle \sigma_0 \rangle \langle \sigma_L \rangle$.
- The connected correlation:

$$\langle \sigma_0 \sigma_L \rangle_c = \langle \sigma_0 \sigma_L \rangle - \langle \sigma_0 \rangle \langle \sigma_L \rangle.$$

Obviously only two of these correlations are linearly independent.

With these two-spins correlations, we can build different squared correlations:

- The total-total correlation at distance L :

$$\begin{aligned} \overline{\langle \sigma_0 \sigma_L \rangle^2}^c &= \frac{2}{\beta^2 z^L} \lim_{n \rightarrow 0} \left[\frac{1}{n(n-1)} \sum_{a \neq b} (T^{L+1})_{ab,ab} \right] = \\ &= \frac{2}{\beta^2 z^L} \left(\frac{3}{2} \lambda_R^{L+1} - \lambda_{L/A}^{L+1} \right. \\ &\quad \left. + (L+1) \lambda_{L/A}^L \frac{\beta^2}{2} (3m_4 - 2m_2) \right). \end{aligned} \quad (6)$$

- The disconnected-disconnected correlation at distance L :

$$\begin{aligned} \overline{\langle \sigma_0 \rangle^2 \langle \sigma_L \rangle^2}^c &= \\ &= \frac{2}{\beta^2 z^L} \lim_{n \rightarrow 0} \left[\frac{1}{n(n-1)(n-2)(n-3)} \sum_{a \neq b \neq c \neq d} (T^{L+1})_{ab,cd} \right] = \\ &= \frac{2}{\beta^2 z^L} \left(\frac{\lambda_R^{L+1}}{2} - \frac{\lambda_{L/A}^{L+1}}{2} \right. \\ &\quad \left. + (L+1) \lambda_{L/A}^L \frac{\beta^2}{2} (3m_4 - 2m_2) \right). \end{aligned} \quad (7)$$

- The total-disconnected correlation at distance L :

$$\begin{aligned} \overline{\langle \sigma_0 \sigma_L \rangle \langle \sigma_0 \rangle \langle \sigma_L \rangle}^c &= \\ &= \frac{2}{\beta^2 z^L} \lim_{n \rightarrow 0} \left[\frac{1}{n(n-1)(n-2)} \sum_{a \neq b \neq c} (T^{L+1})_{ab,ac} \right] = \\ &= \frac{2}{\beta^2 z^L} \left(\frac{3\lambda_R^{L+1}}{4} - \frac{3\lambda_{L/A}^{L+1}}{4} \right. \\ &\quad \left. + (L+1) \lambda_{L/A}^L \frac{\beta^2}{2} (3m_4 - 2m_2) \right). \end{aligned} \quad (8)$$

- The connected-connected correlation at distance L , whose expression can be obtained as a combination of the previous ones:

$$\begin{aligned} \overline{\langle \sigma_0 \sigma_L \rangle_c^2} &= \overline{(\langle \sigma_0 \sigma_L \rangle - \langle \sigma_0 \rangle \langle \sigma_L \rangle)^2} = \overline{\langle \sigma_0 \sigma_L \rangle^2} \\ &\quad - 2 \overline{\langle \sigma_0 \sigma_L \rangle \langle \sigma_0 \rangle \langle \sigma_L \rangle} + \overline{\langle \sigma_0 \rangle^2 \langle \sigma_L \rangle^2} = \\ &= \frac{1}{\beta^2 z^L} \lambda_R^{L+1}. \end{aligned} \quad (9)$$

Also in this case only three of these correlations are linearly independent³.

Others correlations can be readily obtained from the previous one by integration by part. In the SM, we show how to obtain the connected and disconnected bare correlation functions in a cavity approach leading to the same results.

The dominant contribution in the correlation functions. – To build the susceptibility associated with a given correlation function $C(0, L)$ in a Bethe lattice (where C can be one among the correlation functions considered in the previous section), one should sum over all the spins that are at distance L from the spin 0, and then over all the distances L :

$$\chi_C^B \propto \sum_{L=1}^{\infty} \mathcal{N}_L C(0, L), \quad (10)$$

where $\mathcal{N}_L = z(z-1)^{L-1}$ is the number of spins at distance L from a given spin in a Bethe lattice with connectivity z .

³The expression for the connected correlation in the Bethe lattice is obtained in this paper from a large z expansion. However we numerically checked that also for small z eq. (9) is valid substituting z with $z-1$ (that is equivalent in the large z limit).

$$\overline{\langle \sigma_0^a \sigma_0^b \sigma_d^c \sigma_d^d \rangle_{lc}^c} = \left(\frac{\beta^2}{2} \right)^2 \frac{1}{z^{L_0 + (L_1 + 1) + (L_2 + 1) + L_3}} \times \sum_{q,r,s,t} (T^{L_0})_{ab,qr} \left[\sum_{e,f,g,h,l,m,o,p} V_{qr,ef,gh} (T^{L_1})_{ef,lm} (T^{L_2})_{gh,op} V_{st,lm,op} \right] (T^{L_3})_{st,cd}. \quad (13)$$

The SG transition line is commonly associated to the divergence of the SG susceptibility, that is the susceptibility associated to the connected correlation function. Substituting eq. (9) and the expression for \mathcal{N}_L in eq. (10), we discover that the critical line is identified by $\lambda_R = 1$ (see footnote 4). Looking at the eigenvalues, we can numerically check that $\lambda_{L/A} < \lambda_R$. All the bare correlation functions, as shown in the two precedent sections, have a term proportional to λ_R^L , that is thus the dominant one. Recovering the result of standard theory, the critical behavior of all the correlation functions is the same because all depend on the only critical eigenvalue [20]. The susceptibility associated to the different correlations, computed at the critical point, is divergent at the critical line. On the Bethe lattice, this divergence is a consequence of the exponential decay of the correlations multiplied by the exponential numbers of neighbours at a given distance. Until now we have computed χ_C^B on the Bethe lattice. If now we want to use the Bethe approximation to compute the susceptibility in a D -dimensional lattice, following ref. [18] (recall the section “Expanding around the Bethe lattice solution”), we should replace \mathcal{N}_L in eq. (10) with the total number $\mathcal{B}(d, L)$ of non-backtracking paths of length L between two points at distance d on the original lattice, and add a sum over d . For large d and L

$$\mathcal{B}(d, L) \propto (2D - 1)^L |L|^{-D/2} \exp(-|d|^2 / (2L)) \quad (11)$$

implying that at this order the divergence of the susceptibility in a finite-dimensional lattice takes place in correspondence with the divergence in a Bethe lattice with connectivity $z = 2D$.

One spatial loop in the RS phase. – In a Bethe lattice in the thermodynamic limit, the density of loops of finite length vanishes, while spatial loops of finite length are common in finite-dimensional lattices. In ref. [18] a new expansion around the Bethe lattice is performed. As a result, the first correction to the bare propagator computed in the precedent sections comes from one spatial loop. In this section, we will compute this contribution confirming that it is totally equivalent to the first correction computed in the usual field-theoretical loop expansion, as stated in ref. [18]. Following the prescriptions of ref. [18], we construct a spatial loop structure \mathcal{L} , shown in

⁴Indeed one can check that in the limit of $z \rightarrow \infty$ λ_R , as defined in this paper, is deeply related to the replicon eigenvalue λ of eq. (15) from ref. [19]. Using eq. (11) of ref. [19], it can be demonstrated that the relation $(\frac{KT}{J})^2 \lambda = -\lambda_R + 1$ holds. The usual SG line associated with $\lambda = 0$ translates in $\lambda_R = 1$.

fig. 1, formed by two paths of length $L_1 + 1$ and $L_2 + 1$ between the points x and y (the length of the internal paths are defined of length $L_1 + 1$ and $L_2 + 1$ because in this way results are more compact), plus two external legs of length L_0 and L_3 to the external spins σ_0 and σ_d . The rest of the lattice is a Bethe lattice without loops in the thermodynamic limit.

We will compute the correction to the “bare” correlation functions computed in the previous section, that are those connected over the disorder. Analogously to the definition of T , we define the vertex

$$V_{ab,cd,ef}(x) = \overline{\langle \sigma_x^a \sigma_x^b \sigma_x^c \sigma_x^d \sigma_x^e \sigma_x^f \rangle} = \begin{cases} 1, & \text{if three pairs of indices are equal,} \\ m_2, & \text{if two pairs of indices are equal,} \\ m_4, & \text{if one pair of indices is equal,} \\ m_6, & \text{if the indices are all different.} \end{cases}$$

As in the case of T , $a \neq b$, $c \neq d$, $e \neq f$. We write the partition function in the presence of one loop \mathcal{L} and we expand it for small J 's analogously to eq. (1). From it, following the same reasoning of the section “The replica computation of line-correlations”, we obtain the form of the correlation function when a structure \mathcal{L} is present:

$$\overline{\langle \sigma_0^a \sigma_0^b \sigma_d^c \sigma_d^d \rangle_{\mathcal{L}}^c} = \overline{\langle \sigma_0^a \sigma_0^b \sigma_d^c \sigma_d^d \rangle_{L_0, L_1 + 1, L_3}^c} + \overline{\langle \sigma_0^a \sigma_0^b \sigma_d^c \sigma_d^d \rangle_{L_0, L_2 + 1, L_3}^c} + \overline{\langle \sigma_0^a \sigma_0^b \sigma_d^c \sigma_d^d \rangle_{lc}^c}, \quad (12)$$

where the first and second terms turn out to be exactly the “bare” correlations computed as the two paths $L_0 + (L_1 + 1) + L_3$ and $L_0 + (L_2 + 1) + L_3$ were independent. They have, respectively, $L_0 + (L_1 + 1) + L_3$ and $L_0 + (L_2 + 1) + L_3$ couplings, that are the minimal number of couplings to have a non-zero correlation. The last term has $L_0 + (L_1 + 1) + (L_2 + 1) + L_3$ couplings and turns out to be

see eq. (13) above

Thus, in our case, from eq. (12) we see that $\overline{\langle \sigma_0^a \sigma_0^b \sigma_d^c \sigma_d^d \rangle_{lc}^c}$ is exactly the line-connected correlation function, that gives the one-loop correction in the $\frac{1}{M}$ expansion around the Bethe solution. The one-loop contribution takes a very intuitive form.

To compute the explicit form for $\overline{\langle \sigma_0^a \sigma_0^b \sigma_d^c \sigma_d^d \rangle}_{lc}$, we performed sums and products in eq. (13) using Mathematica⁵. At this point we can compute the one-loop contribution to the different correlation functions as in the section “The replica computation of line-correlations”. The whole expressions are reported in the SM. As for the bare term, the dominant terms are those with the highest power of λ_R , that are

$$\begin{aligned} \overline{\langle \sigma_0 \sigma_d \rangle_c}^2|_{lc} &\simeq \overline{\langle \sigma_0 \rangle^2 \langle \sigma_d \rangle^2}|_{lc} \simeq 32\lambda_R^{L_0+L_1+L_2+L_3} \\ &\times [1 + 44m_2^2 + 101m_4^2 + m_4(22 - 90m_6) \\ &- 2m_2(7 + 67m_4 - 30m_6) - 10m_6 + 20m_6^2]. \end{aligned} \quad (14)$$

As explained in the section “The dominant contribution in the correlation functions”, following ref. [18], we can now compute the correction to the susceptibility summing over all the length L_0, L_1, L_2, L_3 , once we have multiplied by the number of non-backtracking walks.

Relation with previous RG studies. – In ref. [14], the authors perform standard RG calculations in $6 - \epsilon$ dimensions for SG with a field. They write the field-theoretic Hamiltonian in the vicinity of the critical line projected on the replicon eigenspace as a function of the order parameter $q_{\alpha\beta}$ as

$$\begin{aligned} H &= \frac{1}{4}r \sum q_{\alpha\beta}^2 + \frac{1}{4} \sum (\nabla q_{\alpha\beta})^2 \\ &- \frac{1}{6}w_1 \sum q_{\alpha\beta} q_{\beta\gamma} q_{\gamma\alpha} - \frac{1}{6}w_2 \sum q_{\alpha\beta}^3 \end{aligned}$$

with r the reduced temperature, and w_1, w_2 being coupling constants. The correlation functions in the momentum space k (projected on the replicon eigenspace) are proportional to $(r + k^2)^{-1}$. Integrating over an infinitesimal shell $e^{-dl} < k < 1$ in the momentum space, they obtain the recursion relation for r :

$$\frac{dr}{dl} = (2 - \eta)r - \frac{K_d}{(1+r)^2}(4w_1^2 - 16w_1w_2 + 11w_2^2), \quad (15)$$

with $\eta = \frac{1}{3}K_d(4w_1^2 - 16w_1w_2 + 11w_2^2)$ and K_d the usual geometrical factor, together with analogous recursion relations for w_1 and w_2 . The first term in eq. (15) is the contribution connected to a renormalization of the critical temperature, and that we can compute in our approach computing the correction given by a spatial tadpole structure. The second term is the one coming from the non-trivial loop.

Following refs. [21,22], in a Bethe lattice in finite and infinite connectivity, as well as in the fully connected model, $w_1 \propto 1 - 3m_2 + 3m_4 - m_6$ and $w_2 \propto 2m_2 - 4m_4 + 2m_6$. Inserting these expressions in the loop term of eq. (15),

⁵The same results can be concluded from eq. (62) of ref. [15] after a suitable correspondence between quantities like the propagators and 3-point vertex.

we obtain

$$\begin{aligned} 4w_1 - 16w_1w_2 + 11w_2^2 &\propto 4 - 56m_2 + 176m_2^2 + 88m_4 \\ &- 536m_2m_4 + 404m_4^2 - 40m_6 \\ &+ 240m_2m_6 - 360m_4m_6 + 80m_6^2, \end{aligned}$$

that is exactly proportional to the coefficient of the dominant term for the spatial-loop correction to the connected correlation function eq. (14) (see footnote ⁶). Our approach permits to find the same results in a clearer, simpler and more physically intuitive way.

Conclusions and perspectives. – In ref. [19] a new expansion is introduced around the Bethe lattice. This expansion can be applied to all the models that can be defined on a Bethe lattice. It is supposed to give the same results as standard perturbation theory for fully connected models when the physics of fully connected and finite connectivity MF models is the same. We test the new expansion of ref. [18] for the first time in the case of the spin-glass model at finite temperature and with an external field in the limit of high connectivity. We analytically find the same results of standard RG [14], confirming the validity of the new M -layer topological expansion.

The new method has, however, the advantage that the equations are physically very intuitive: the bare correlations are just the correlation on a Bethe lattice (*i.e.*, on a line), and the first-order correction to the correlation function is just the value of the correlation function computed on a spatial loop (finite loops are absent in the Bethe lattice and present in finite-dimensional systems) once the contributions of the two lines forming the loop, considered as independent graphs, are subtracted. Even if for the case of the SG in a field in the limit $z \rightarrow \infty$ and finite temperature the results of the new expansion add nothing to what already known about the SGs in finite dimensions, there are cases in which things should be different.

It was already underlined how the Bethe lattice is more similar to finite-dimensional systems [23] with respect to the FC version for different disordered spin models. In the particular case of SG in a field, the critical line on the FC model tends to infinite field when the temperature goes to zero, while in the Bethe lattice it ends at a finite field h_c at $T = 0$ [21]. If the critical point for SGs in field in finite dimension is a zero-temperature one, as supposed by some authors [5,6], it is crucial to perform an expansion around the Bethe solution, instead of around FC model, since the latter model is always in the SG phase at $T = 0$. This paper is a first step in this direction. The application of the topological expansion to the SG model in a field at $T = 0$ and finite connectivity is at the moment under study. It could be done according to lines similar to what done in the case of the random field Ising model [24].

⁶Please note that the replicon contribution, that is the one multiplied by the coefficient b_4 , that corresponds to the one found by Bray and Roberts, is the dominant one for the correlation functions. In our approach, we obtain also the sub-dominant corrections coming from the other sectors, as in refs. [15,16].

We thank TOMMASO RIZZO for very useful discussions.

REFERENCES

- [1] SHERRINGTON D. and KIRKPATRICK S., *Phys. Rev. Lett.*, **35** (1975) 1792.
- [2] PARISI G., *J. Phys. A*, **13** (1980) L115; **13** (1980) 1101.
- [3] PANCHENKO D., *Curr. Dev. Math.*, **2014** (2016) 231.
- [4] MOORE M. A. and BRAY A. J., *Phys. Rev. B*, **83** (2011) 224408.
- [5] PARISI G. and TEMESVARI T., *Nucl. Phys. B*, **858** (2012) 293.
- [6] ANGELINI M. C. and BIROLI G., *Phys. Rev. Lett.*, **114** (2015) 095701.
- [7] CHARBONNEAU P. and YAIDA S., *Phys. Rev. Lett.*, **118** (2017) 215701.
- [8] LARSON D., KATZGRABER H. G., MOORE M. A. and YOUNG A. P., *Phys. Rev. B*, **87** (2013) 024414.
- [9] KATZGRABER H. G., LARSON D. and YOUNG A. P., *Phys. Rev. Lett.*, **102** (2009) 177205.
- [10] LEUZZI L., PARISI G., RICCI-TERSENGHI F. and RUIZ-LORENZO J. J., *Phys. Rev. Lett.*, **103** (2009) 267201.
- [11] JANUS COLLABORATION, *Phys. Rev. E*, **89** (2014) 032140; *J. Stat. Mech.* (2014) P05014.
- [12] MATTSSON J., JONSSON T., NORDBLAD P., ARUGA KATORI H. and ITO A., *Phys. Rev. Lett.*, **74** (1995) 4305.
- [13] HARRIS A. B., LUBENSKY T. C. and CHEN J.-H., *Phys. Rev. Lett.*, **36** (1976) 415.
- [14] BRAY A. J. and ROBERTS S. A., *J. Phys. C*, **13** (1980) 5405.
- [15] TEMESVARI T., DE DOMINICIS C. and PIMENTEL I. R., *Eur. Phys. J. B*, **25** (2002) 361.
- [16] PIMENTEL I. R., TEMESVARI T. and DE DOMINICIS C., *Phys. Rev. B*, **65** (2002) 224420.
- [17] TEMESVARI T., *Phys. Rev. B*, **96** (2017) 024411.
- [18] ALTIERI A., ANGELINI M. C., LUCIBELLO C., PARISI G., RICCI-TERSENGHI F. and RIZZO T., *J. Stat. Mech.* (2017) 113303.
- [19] DE ALMEIDA J. R. L. and THOULESS D. J., *J. Phys. A*, **11** (1978) 983.
- [20] DE DOMINICIS C. and GIARDINA I., *Random Fields and Spin Glasses: A Field Theory Approach* (Cambridge University Press) 2006.
- [21] PARISI G., RICCI-TERSENGHI F. and RIZZO T., *J. Stat. Mech.* (2014) P04013; (2015) E03001.
- [22] SOMPOLINSKY H. and ZIPPELIUS A., *Phys. Rev. B*, **25** (1982) 6860.
- [23] CAMMAROTA C., BIROLI G., TARZIA M. and TARJUS G., *Phys. Rev. B*, **87** (2013) 064202.
- [24] ANGELINI M. C. *et al.*, in preparation.